

Lysozyme activity of the *Ruminococcus champanellensis* cellulosome

Sarah Morais,¹ Darrell W. Cockburn,²
Yonit Ben-David,¹ Nicole M. Koropatkin,²
Eric C. Martens,² Sylvia H. Duncan,³ Harry J. Flint,³
Itzhak Mizrahi⁴ and Edward A. Bayer^{1*}

¹Biomolecular Sciences Department, The Weizmann Institute of Science, Rehovot, Israel.

²Department of Microbiology and Immunology, University of Michigan Medical School, Ann Arbor, MI 48109, USA.

³Microbiology Group, Rowett Institute of Nutrition and Health, University of Aberdeen, Aberdeen, UK.

⁴The Department of Life Sciences & the National Institute for Biotechnology in the Negev, Ben-Gurion University of the Negev, Beer-Sheva 84105, Israel.

Summary

Ruminococcus champanellensis is a keystone species in the human gut that produces an intricate cellulosome system of various architectures. A variety of cellulosomal enzymes have been identified, which exhibit a range of hydrolytic activities on lignocellulosic substrates. We describe herein a unique *R. champanellensis* scaffoldin, ScaK, which is expressed during growth on cellobiose and comprises a cohesin module and a family 25 glycoside hydrolase (GH25). The GH25 is non-autolytic and exhibits lysozyme-mediated lytic activity against several bacterial species. Despite the narrow acidic pH curve, the enzyme is active along a temperature range from 2 to 85°C and is stable at very high temperatures for extended incubation periods. The ScaK cohesin was shown to bind selectively to the dockerin of a monovalent scaffoldin (ScaG), thus enabling formation of a cell-free cellulosome, whereby ScaG interacts with a divalent scaffoldin (ScaA) that bears the enzymes either directly or through additional monovalent scaffoldins (ScaC and ScaD). The ScaK cohesin also interacts with the dockerin of a protein comprising multiple Fn3 domains that can potentially

promote adhesion to carbohydrates and the bacterial cell surface. A cell-free cellulosomal GH25 lysozyme may provide a bacterial strategy to both hydrolyze lignocellulose and repel eventual food competitors and/or cheaters.

Introduction

Ruminococcus champanellensis is the sole known human gut bacterium able to degrade crystalline cellulose (Chassard *et al.*, 2012). It produces a cellulosome that has been characterized recently (Ben David *et al.*, 2015; Morais *et al.*, 2016). Cellulosomes are large-molecular-weight enzymatic complexes that represent an extremely efficient strategy for cellulose and hemicellulose degradation (Bayer *et al.*, 2004; 2007; Himmel *et al.*, 2010). The strong intermodular calcium-dependent cohesin/dockerin interaction drives the assembly between the cellulosomal enzymes and the central non-catalytic, integrating subunit, the scaffoldin, to form the mature complex (Yaron *et al.*, 1995; Lytle *et al.*, 1996). The *R. champanellensis* genome contains 12 scaffoldins with various molecular arrangements and specificities (Fig. 1). The largest cellulosome that could be assembled by *R. champanellensis* would be composed of an anchoring scaffoldin, ScaE, comprising a sortase motif and a cohesin that can interact with the dockerin of a second scaffoldin, ScaB. The latter contains seven cohesin modules, three of which interact either directly with dockerin-bearing enzymes, whereas the remainder can interact either with dockerin-bearing enzymes or with a third dockerin-bearing scaffoldin, ScaA. ScaA also contains two cohesin modules that interact either directly with dockerin-bearing enzymes or indirectly via monovalent scaffoldins (ScaC and ScaD) that play a role of molecular adaptors, thereby modulating the enzymatic composition of the cellulosome (Rincon *et al.*, 2004; Ben David *et al.*, 2015). The entire complex would thus contain a maximum of 11 enzymes. *R. champanellensis* contains a total of 65 dockerin-bearing proteins, 8 scaffoldin-borne dockerins (Ben David *et al.*, 2015), 25 recently characterized glycoside hydrolases (Morais *et al.*, 2016) and 31 additional dockerin-containing proteins. The *R. champanellensis* cellulosome thus presents a finely-tuned cohesin/dockerin recognition system that enables

Received 6 March, 2016; accepted 16 August, 2016. *For correspondence. Email: ed.bayer@weizmann.ac.il; Tel: (+972) 8 934 2373; Fax: (+972) 8 934 4118.




Fig. 1. Schematic representation of the cohesin-bearing scaffoldin proteins in *R. champanellensis* based on the respective genome sequences.

SGNH, hydrolase-type esterase domain (IPR013830); GH25, a putative GH25-family module sharing similarity to lysozyme. Specific interactions according to sequence alignment and biochemically characterized cohesin/dockerin interactions are color-coded. Shaded colors indicate that the designated cohesins or dockerins exhibit more selective interactions with only some of their Group 1 or Group 2 counterparts.

regulated assembly of an elaborate cellulosomal organization (Morais *et al.*, 2016).

Unlike common scaffoldins that are non-enzymatic subunits, the ScaK scaffoldin carries an enzyme that is associated with the metabolism of cellular structural components, i.e., the peptidoglycan. ScaK is composed of a cohesin module (definitive of the scaffoldins) at its N-terminus and a GH25 catalytic domain in its C-terminal region. GH25 enzymes are retaining glycoside hydrolases that cleave the β -1,4-glycosidic bond between *N*-acetylmuramic acid (NAM) and *N*-acetylglucosamine (NAG) in the carbohydrate backbone of bacterial peptidoglycan (i.e., lysozymes). Two main biological roles can be attributed to bacterial GH25 enzymes: the autolysis, the re-modeling of peptidoglycan in cellular growth processes, and the dissemination of phage progeny towards the end of the phage lytic cycle via lysis of the bacterial cell. These specialized hydrolases can also create enlarged pores in the bacterial peptidoglycan for the assembly of large trans-envelope complexes (e.g., pili, flagella, secretion systems) (Vollmer *et al.*, 2008).

The genome of *Ruminococcus* sp. CAG:379 strain, closely related to *R. champanellensis*, also contains a gene encoding a protein homologous to ScaK with a similar modular arrangement (cohesin and GH25 modules). In additional *Ruminococcus* species, such as *R. bicirculans* and *R. flavefaciens* that carry multiple GH25 genes, the GH25 module is not accompanied by a cohesin module. ScaK also has 30% identity and 49% similarity with the GH25 module of LytC, an autolysin from *Streptococcus pneumoniae*, which is involved in a three-component mechanism for the lysis of sister cells non-competent for natural genetic transformation (Claverys *et al.*, 2007; Monterroso *et al.*, 2008; Eldholm *et al.*, 2009). The fact that the ScaK GH25 module is connected to a cellulosomal element raises the question of possible lysozyme activity associated with the cellulosome that could provide a novel strategy that would prove beneficial for the bacterial cell.

In our previous report (Ben David *et al.*, 2015), the ScaK cohesin was expressed as a CBM-fused cohesin (carbohydrate binding module) but was not found to bind any of the dockerin partners tested. In the present study, we cloned and expressed the ScaK scaffoldin as an intact wild-type protein, in an attempt to reveal potential cohesin affinity partners and to study the GH25 activity.

Results

Scaffoldin production and functionality of their cohesin modules

ScaK, ScaG and ScaF scaffoldins (Fig. 1) were produced in *Escherichia coli*, purified on Ni-beads, and their estimated molecular weights were in good agreement with their calculated molecular masses (see legend to Supporting Information Fig. S1).

In order to understand the architectural context in which ScaK is positioned within the cellulosome, the dockerin specificities of the ScaK cohesin was examined by an affinity ELISA approach using the representative dockerin-bearing fusion proteins listed in Table 1. For this purpose, 12 dockerin modules were fused to a xylanase tag (xylanase T6 from *Geobacillus stearothermophilus*) that promotes solubility and expression of the dockerin and serves as a recognition tag for the primary antibody in the ELISA procedure (Barak *et al.*, 2005). The resultant xylanase-dockerin (Xyn-Doc) fusion proteins are listed in Table 1. All dockerins from the *R. champanellensis* scaffoldins (eight in total) were tested, since they represent the main backbones of its cellulosome structures. The four additional selected dockerins belonged to the three dockerin groups (Groups 1, 2 and 3–4) of cohesin–dockerin interactions in this bacterium, as previously defined using bioinformatic-based criteria (Ben David *et al.*, 2015; Morais *et al.*, 2016). ScaK, expressed as a full-length protein (lacking the signal peptide), was able to interact with the

Table 1. List of the *R. champanellensis* Xyn-fused dockerin proteins used in this study. Name and modular architecture of the original scaffoldin or protein are given.

Fused dockerin	Parent scaffoldin/protein	Modular architecture	Dockerin group
Xyn-DocA	ScaA	SIGN X Coh Coh Doc	2
Xyn-DocB	ScaB	SIGN Coh Coh Coh Coh Coh Coh Coh Coh X Doc	1
Xyn-DocC	ScaC	SIGN Coh UNK Doc	2
Xyn-DocD	ScaD	SIGN Coh Doc	2
Xyn-DocF	ScaF	SIGN Coh Doc	1
Xyn-DocG	ScaG	SIGN Coh Doc	1
Xyn-DocH	ScaH	SIGN SGNH Coh Doc	1
Xyn-DocJ	ScaJ	SIGN Coh Coh Coh Doc	1
Xyn-Doc3939	gi:291543939	SIGN Fn3 PKD Fn3 Fn3 Fn3 PKD Doc	1
Xyn-Doc48	Cel48A	SIGN GH48A Doc	2
Xyn-Doc9B	GH9B	SIGN CBM4 Fn3 GH9B Doc GH16A	3–4
Xyn-Doc3550	gi:291543550	UNK Doc	3–4

The numbers for proteins 3550 and 3939 refer to the last four digits of the respective full GI number (i.e., 29154XXXX).


Abbreviations: Xyn, XynT6 from the *G. stearothermophilus*; SIGN, signal peptide; Doc, dockerin; Coh, cohesin; GH, glycoside hydrolase; SGNH, lipases or esterases; Fn3, fibronectin type III; PKD, polycystic kidney disease; UNK, X, unknown.

ScaG and protein 3939 dockerins (Fig. 2A), in contrast to the previous report (Ben David *et al.*, 2015), where the cohesin module was expressed alone, in the absence of the GH25 module.

ScaG is a scaffoldin that comprises only a cohesin and a dockerin (Fig. 1). In a previous report (Ben David *et al.*, 2015), the cohesin and dockerin modules of ScaG were expressed separately, the dockerin demonstrated binding activity for the first cohesin of ScaJ and ScaE, but the cohesin of ScaG failed to interact with any of the dockerin counterparts and was termed inactive in that study. We produced here the full ScaG protein length in an attempt to reveal the binding partner of its cohesin. Using this approach, the ScaG cohesin was able to bind to the ScaA dockerin (Figs. 1 and 2B), suggesting an adaptor role for

this scaffoldin between ScaK and the enzyme-bearing scaffoldin ScaA. A scheme illustrating the interactions between ScaK and the cellulosomal elements is presented in Fig. 3. Consequently, the ScaG cohesin is now considered a Group-2 cohesin, and the ScaK cohesin and dockerins of ScaA and Prot3939 belong to the Group-1 interacting modules.

ScaF exhibits the same modular architecture as ScaG (Fig. 1), but with a longer linker region at the N-terminus of the protein. Similar to ScaG, when the cohesin and dockerin modules were expressed separately, only the dockerin was active and exhibited similar binding abilities as the ScaG dockerin (Ben David *et al.*, 2015). Similar to ScaK and ScaG, the expression of the full-length scaffoldin revived the cohesin activity, and specific interaction with

**Fig. 2.** Newly discovered dockerin-binding profiles of *R. champanellensis* ScaK and ScaG cohesins measured by affinity ELISA.

A. ELISA experiments demonstrating different interaction specificities between the ScaK cohesin and selected scaffoldin- and enzyme-borne dockerins. ScaK interacted with DocG and Doc3939.

B. ELISA experiments demonstrating different interaction specificities between the ScaG cohesin and selected scaffoldin- and enzyme-borne dockerins. ScaG interacted exclusively with DocA. Error bars indicate the standard deviation from the mean of triplicate samples (ELISA). The experiments were performed three times.




Fig. 3. Proposed cell-free cellulosome complexes involving ScaK in *R. champanellensis*. Different types of cohesin-dockerin interactions are color-coded. The ScaK cohesin binds selectively to the dockerins of ScaG and Prot3939 and not to other Group 1 dockerins. Similarly, the ScaG cohesin appears to be very selective in its binding to the ScaA dockerin, and fails to bind other Group 2 dockerins.

the ScaG dockerin was observed (Supporting Information Fig. S2). As ScaG and ScaF dockerins exhibited the same binding affinities and both can bind directly to the ScaE cohesin, it is not clear why the bacterium would produce and assemble an additional adaptor scaffoldin (i.e., ScaF) to mediate between ScaG and ScaE (via ScaJ) (Supporting Information Fig. S3). In this context, ScaF could perhaps serve as an extender to increase the overall length of the cellulosome in order to avoid steric hindrance or to access more distant substrates.

Lysozyme activity

The enzymatic activity was monitored by decrease in turbidity of the cultures. The lysozyme module of ScaK was active on *E. coli* cells between pH 3.5 and 5.5 with a pH optimum of 5 (Fig. 4A). The activity of the enzyme was lost completely at pH 6 and higher. The protein was stable between 30 and 60°C for 48 h. At 70°C, the lysozyme retained its full activity after 3 h of incubation and was 54% active after 24 h of incubation. After 48 h of incubation at this temperature, the enzyme was still 30% active. At 80°C, the enzyme was more rapidly degraded, but was still 20% active after 48 h of incubation (Fig. 4B). ScaK was active on *E. coli* cells from 2 to 45°C. Enzymatic activity increased with the temperature as monitored by the decrease in turbidity and viable cell count (Fig. 4C). In addition, the enzyme was active from 2 to 85°C on purified peptidoglycans from *Bacillus subtilis* and *Staphylococcus aureus* along the entire temperature range with moderate enhancement of activity with increasing temperature (Fig. 4D).

The lytic activity of ScaK was tested against a large number of strains that colonize the human gut and additional bacteria (Table 2). The lysozyme was found to be inactive

against the parent *R. champanellensis* cells but active against several *Lactobacillus* strains, *E. coli*, *Enterococcus faecalis*, *Listeria monocytogenes* and *S. pneumoniae*. *Clostridium difficile* was degraded after long incubation periods. Growth of *E. coli* and *Lactobacillus plantarum* was inhibited by the presence of the lysozyme, as demonstrated by inhibition zones around disks containing the enzyme during growth on plates (Supporting Information Fig. S4).

The potential protective role of the bacterial capsule against bacterial cell lysis was tested by incubating the lysozyme with *E. coli* strains producing a capsule (5911 and Nissle strains), and no lytic action was observed. In addition, *Bacteroides thetaiotaomicron*, was tested using both the wild-type strain and a mutant lacking any capsule (Rogers *et al.*, 2013), and in both cases no lysis was observed (Table 2).

Expression of ScaK by *R. champanellensis*

ScaK was detected by proteomic analysis in supernatant fluids of *R. champanellensis* cultures grown on cellobiose as the sole carbon source (score of 9.45 and % coverage of 9.23, p -value < 0.01), suggesting that it is indeed incorporated in the cellulosomal complex and participates in bacterial lignocellulose degradation as a possible inhibitor of other competing or deleterious bacteria which inhabit the human gastrointestinal tract.

Discussion

The ScaK scaffoldin of *R. champanellensis* contains the first cellulosome-associated GH25 known to date. The enzyme is active at extremes of temperature and stable for long incubation periods at elevated temperatures. These characteristics could therefore render the ScaK enzyme a




Fig. 4. Characterization of recombinant *R. champanellensis* ScaK.

A. Effect of pH on lysozyme activity after 15-min incubation.

B. Thermostability of ScaK-derived lysozyme activity at different temperatures, following the heat-shock, the enzymatic activity was measured at pH 5 after 30 min incubation at 37°C.

C. Effect of temperature on lysozyme activity on *E. coli* cells at pH 5 after 30 min incubation, solid line representing the decrease in OD at 600 nm and dashed line, the viability of the cells (in cfu ml^{-1}) with the control counting 1.6×10^{12} cfu ml^{-1} .

D. Effect of temperature on lysozyme activity on the peptidoglycans of *B. subtilis* (blue line) and *S. aureus* (red line) at pH 5 after 30-min incubation. Enzymatic reactions, performed using the modified turbidimetric method were repeated in triplicate, and standard deviations are indicated.

suitable candidate for industrial applications, for example, as an antimicrobial agent in the food industry (Datta *et al.*, 2008; Callewaert *et al.*, 2011; Aminlari *et al.*, 2014) and may also be considered as a therapeutic agent in humans (Pastagia *et al.*, 2013).

Curiously, ruminococci seem to carry numerous non-cellulosomal GH25 genes – nine in *R. flavefaciens* FD1, five in *R. bicirculans* 80/3, five in *R. albus* 7, one in *R. bromii* and three in *R. champanellensis* 18P13 (in addition to the ScaK gene). Since genomic analysis has indicated that prophage DNA (encoding lysozymes) is rather common in this group of bacteria (Berg Miller *et al.*, 2012; Wegmann *et al.*, 2014), we analyzed the flanking regions of the ScaK gene for the presence of phage genes, such as holins (Supporting Information Table 1). In the absence of such genes, bacteriophage origin would thus be improbable. Therefore, the association of the cohesin module and a GH25 gene only with the *R. champanellensis* GH25 raises the question of a biological role for this scaffoldin in its cellulosomal complex.

The production of ScaK, ScaG and ScaF as intact proteins revived the binding activity of their lone cohesins

towards the specified dockerins, suggesting that the functional status and/or structural stability of the cohesin modules of these scaffoldins is dependent on the full protein architecture. We could thus elucidate their dockerin specificities and therefore complete the proposed architecture of the *R. champanellensis* cellulosome system. The results indicate that cell-free cellulosomes can be formed between ScaK that interacts with ScaG, which in turn interacts with ScaA that harbors the enzymes either directly or through ScaC and ScaD adaptors. This type of cellulosome could thus be targeting plant-derived lignocellulosic materials located at a distance from the cell.

The ScaK cohesin was also found to interact with the dockerin of protein 3939. This very large protein has an estimated molecular weight of 308 kDa, and annotation of its sequence has revealed five predicted Fn3 (fibronectin type III) and two PKD (polycystic kidney disease) domains. The Fn3 domains are relatively common components in cellulolytic bacteria, yet their function is not completely understood. Several studies suggested that these domains can mediate protein assembly, adhesion to carbohydrate

Table 2. Bacterial strains tested for ScaK lytic action.

Bacterial phyla/(family)/species/strain	Gram	OD _{T=0}	OD _{T=1h}	ScaK lytic action
Actinobacteria				
(Bifidobacteriaceae)				
<i>Bifidobacterium adolescentis</i> L2-32	+	0.132	0.124	–
(Coriobacteriaceae)				
<i>Collinsella aerofaciens</i> DSM 3979	+	0.094	0.095	–
Firmicutes				
(Bacillaceae)				
<i>Bacillus subtilis</i> NCIB 3610	+	1.200	1.195	–
<i>Bacillus subtilis</i> PY79	+	1.220	1.199	–
(Enterococcaceae)				
<i>Enterococcus faecalis</i> JH2-2	+	0.299	0.210	+
(Lachnospiraceae)				
<i>Eubacterium hallii</i> L2-7	+	0.085	0.074	–
<i>Eubacterium rectale</i> A1-86 (DSM17629)	+	0.925	0.884	–
<i>Anaerostipes hadrus</i> SS2/1	+	1.100	1.029	–
<i>Lachnospiraceae</i> sp. nov. M62/1	+	0.085	0.088	–
<i>Blautia obeum</i> A2-162	+	1.030	0.984	–
<i>Roseburia faecis</i> M72/1	+	0.862	0.856	–
(Lactobacillaceae)				
<i>Lactobacillus plantarum</i> WCFS1	+	1.840	0.368	+
<i>Lactobacillus pentosus</i> DSM 20314	+	1.560	0.267	+
<i>Lactobacillus reuteri</i> Ca6	+	0.210	0.214	–
<i>Lactobacillus rhamnosus</i> GG	+	1.952	0.341	+
(Listeriaceae)				
<i>Listeria monocytogenes</i> 10403S	+	1.194	0.467	+
(Peptostreptococcaceae)				
<i>Clostridium difficile</i> VPI 10463	+	0.470	0.418 (T=4h)	±
(Ruminococcaceae)				
<i>Ruminococcus champanellensis</i> 18P13	+	1.130	1.259	–
<i>Ruminococcus bicirculans</i> 80/3	+	0.459	0.444	–
<i>Ruminococcus bromii</i> L2-63	+	0.382	0.361	–
<i>Eubacterium siraeum</i> 70/3	+	0.437	0.422	–
<i>Faecalibacterium prausnitzii</i> A2-165	+	0.336	0.335	–
<i>Ruminococcus flavefaciens</i> 17	+	0.331	0.330	–
<i>Ruminococcus albus</i> SY3	+	0.689	0.665	–
<i>Clostridium clariflavum</i> DSM 19732	+	1.345	1.322	–
<i>Clostridium thermocellum</i> DSM 1313	+	1.202	1.197	–
(Streptococcaceae)				
<i>Lactococcus lactis</i> MG5267	+	1.500	1.501	–
<i>Lactococcus lactis</i> Z3000	+	1.500	1.499	–
<i>Streptococcus gordonii</i> DL-1	+	0.109	0.100	–
<i>Streptococcus pneumoniae</i> R6	+	0.480	0.051	+
Bacteroidetes				
(Bacteroidaceae)				
<i>Bacteroides thetaiotaomicron</i> ATCC 29148	–	1.573	1.559	–
<i>Bacteroides thetaiotaomicron</i> ΔCPS	–	1.567	1.509	–
Proteobacteria				
(Enterobacteriaceae)				
<i>Escherichia coli</i> BL21	–	1.462	0.206	+
<i>Escherichia coli</i> 5911 (K12)	–	1.500	0.190	+
<i>Escherichia coli</i> 5911 (K12) ^a	–	1.423	1.360	–
<i>Escherichia coli</i> Nissle	–	1.692	1.666	–
(Moraxellaceae)				
<i>Acinetobacter baumannii</i> ATCC 17978	–	0.689	0.648	–
(Vibrionaceae)				
<i>Vibrio cholera</i> TRH7000	–	0.656	0.619	–

a. Grown with sucrose.

Enzymatic activity was monitored by decrease in turbidimetry in cell cultures at 600 nm, from the beginning OD_{T=0} to the end of the reaction OD_{T=1h}. Classification of the Firmicutes based on the work by Ludwig *et al.* (2009).

substrates and/or bacterial cell surfaces (which is in accordance with lysozyme activity). Alternatively, Fn3 domains have been suggested to play a role as flexible peptide linkers or to facilitate the solubility of large protein complexes (Devillard *et al.*, 2004; Alahuhta *et al.*, 2010). In addition, PKD domains have also been found to be involved in protein–protein and/or protein–carbohydrate interactions (Lohning *et al.*, 1996). Therefore, we can assume that protein 3939 may have an important, but largely undefined role in carbohydrate degradation, which would benefit the bacterium in general and would justify the expression and secretion of such a large protein. The dockerin of protein 3939 was previously reported to also interact with ScaE that is anchored to the bacterial cell wall (Ben David *et al.*, 2015).

It is also of note that other putative lysozymes can be encoded by GH23-, GH24- and GH73-containing genes in other bacterial genomes. Interestingly, the sole dockerin of the *R. bicirculans* genome was associated with a GH73 catalytic domain (Wegmann *et al.*, 2014), suggesting a similar mechanism and role for the lysozyme in this bacterium.

It is plausible that the GH25 lysozyme has a defensive role against other carbohydrate-degrading bacterial competitors by targeting and effecting the lysis of bacterial cells in the vicinity that would compete for the enzymatic degradation products. In nature, *R. champanellensis* and other specialized fiber-degrading bacteria are prone to exploitation by cheating microbes that utilize the fiber-released soluble sugars, taking advantage of the bacterial fiber-degrading machineries carried without paying the fitness cost (Berlemont and Martiny, 2013). It is therefore reasonable to assume that the fiber degraders will develop defense mechanisms protecting them from such exploitation. In this sense, the GH25 lysozyme exhibits an efficient selective tactic, as it does not affect the parent *R. champanellensis* cell itself, but lyses other gut microbial cells (Table 2).

The bacterial spectrum of susceptibility to ScaK is quite heterogeneous, whereby some Gram-positive (*Lactobacillus* species, *Enterococcus faecalis*, *Listeria monocytogenes*, *S. pneumoniae* and *Clostridium difficile*) and Gram-negative (*E. coli*) bacteria were lysed. Our results initially suggested that ScaK specifically degraded peptidoglycans that contain meso-diaminopimelate (*m*-DAP) residues in the peptidoglycan structure (Schleifer and Kandler, 1972; Humann and Lenz, 2009), since all the strains that were herein susceptible to ScaK lytic action contained *m*-DAP. However, since the isolated purified peptidoglycan of *S. aureus*, which contains D-alanine residue and not *m*-DAP, was degraded, this hypothesis was thus invalidated.

Bacterial defense mechanisms against lysozyme activity are diverse and include the production of enzyme inhibitors or modification of the peptidoglycan (Callewaert *et al.*, 2012), as well as the production of bacterial capsules

(Fouet and Mesnage, 2002). We thus investigated the role of the bacterial capsule as a possible protection against the lysozyme action by testing *E. coli* strains with and without capsules. In the case of the two encapsulated strains of *E. coli*, i.e., Nissle and 5911, the capsule indeed conferred resistance to ScaK lytic action. It should also be noted that capsule expression is highly controlled and regulated (Torres-Cabassa and Gottesman, 1987; Gottesman and Stout, 1991; Sledjeski and Gottesman, 1996) and that four different types of bacterial capsules are described to date in *E. coli* (Whitfield and Roberts, 1999). The strains tested herein belong to Group 2 capsules (capsular gene K5 and K12 for Nissle and 5911 strains, respectively). Therefore, other groups of capsules may confer resistance to the lytic action of ScaK or not, depending also on the growth conditions and extent of capsule expression. We also investigated the capsular protective role with two different strains of *Bacteroides thetaiotaomicron* (Shah, 1992). The two strains tested were the wild-type strain and a mutant lacking any capsule. In both cases, no lysis was observed. Consequently, together with the non-susceptible Gram-negative bacteria tested herein, the outer membrane may act as an impermeable barrier for small molecules thereby protecting the peptidoglycan layer from the lytic action of the lysin (Briers and Lavigne, 2015). Nevertheless, some endolysins have been reported to cross the outer membrane and lyse the bacteria (Lai *et al.*, 2011; Lood *et al.*, 2015). This ability could reflect highly positively charged N- or C-terminal domains in their protein sequence, which enable the lysins to bind to the anionic outer membrane and access their peptidoglycan substrate (Lai *et al.*, 2011). In our case, we could not identify a similar domain in ScaK that could account for the lysis of the Gram-negative acapsulated strains of *E. coli* (*BL21* and *5911*).

In conclusion, we identified the presence of a non-autolytic lysozyme in the *R. champanellensis* cellulosome system. The production of this cell-free cellulosome would represent a strategy to hydrolyze lignocellulose while repelling eventual food competitors or cheaters. The fact that this lysozyme activity is associated with glycoside hydrolases in a single cell-free cellulosome complex suggests a broader role for cellulosomal complexes that would not be restricted to plant cell wall deconstruction.

Experimental procedures

Cloning

Scaffolds ScaK, ScaG and ScaF were cloned from *R. champanellensis* genomic DNA using Phusion High Fidelity DNA polymerase F530-S (New England Biolabs) and the following primers. For ScaK amplification, the primer pair 5'-ttactaCCA TGGcacaccatcaccatcaccatgcagatcagactgtacagac-3' and 5'-ttactaCTCGAGttaaaccattaaatccgt-3' was used, for ScaG, 5'-tact

gaCCATGGcacaccatcaccatcaccatcagaccatgcagccggcgccg-3' and 5'-tacttaCTCGAGtcaaccgagcaggtcatccc-3' and for ScaF 5'-tactgaCCATGGcacaccatcaccatcaccatcaccatcagaccatgcagccggcgccg-3' and 5'-tacgatCTCGAGtcaccattgctggattcggatc-3'. NcoI and XhoI (restriction sites in uppercase). FastDigest enzymes (Thermo scientific, USA) were incubated with the PCR products before their ligation into linearized pET28a using T4 DNA ligase (Fermentas UAB, Vilnius, Lithuania). PCR products were purified using a HiYield™ Gel/PCR Fragments Extraction Kit (Real Biotech Corporation, RBC, Taiwan), and plasmids were extracted using Qiagen miniprep kit (The Netherlands). Competent *E. coli* XL1 cells were used for plasmid transformation.

The full list of fused dockerins used in this article is given in Table 1.

Recombinant protein expression and purification

E. coli BL21 (DE3) cells were transformed with the desired plasmid and plated onto LB-kanamycin plates. The cells producing ScaK, ScaG and ScaF were grown in 500 ml LB (Luria Broth) and 2 mM CaCl₂ at 37°C until $A_{600} \approx 0.8$ –1. The cells were induced by adding 0.1 mM (final concentration) of isopropyl-1-thio- β -D-galactoside (IPTG) (Fermentas UAB Vilnius, Lithuania), and cell growth was continued at 16°C overnight. Cells were harvested by centrifugation at 5000 rpm for 5 min. Pelleted cells were resuspended in 1 ml TBS (Tris-buffered saline, 137 mM NaCl, 2.7 mM KCL, 25 mM Tris-HCl, pH 7.4). Xyn-Doc proteins were expressed as described previously (Ben David *et al.*, 2015; Morais *et al.*, 2016).

His-tagged proteins (scaffoldins and Xyn-Doc) were purified on a Ni-NTA column (Qiagen), as reported earlier (Caspi *et al.*, 2006).

Purity of the recombinant proteins was tested by SDS-PAGE on 10% acrylamide gels, and protein concentration was estimated by absorbance (280 nm), based on the known amino acid composition of the protein, using the ProtParam tool (Gasteiger *et al.*, 2005). Proteins were stored in 50% (vol/vol) glycerol at -20°C .

Affinity-based ELISA

The matching fusion-protein procedure by Barak *et al.* (Barak *et al.*, 2005; Caspi *et al.*, 2006) was followed to determine cohesin–dockerin specificity of interaction. Scaffoldins were immobilized on the plate at a concentration of $1 \mu\text{g ml}^{-1}$ (100 $\mu\text{l/well}$) in 0.1 M sodium carbonate (pH 9) and incubated at 4°C overnight. The following steps were performed at room temperature for 1 h with all reagents at a volume of 100 $\mu\text{l/well}$ with a washing step (300 $\mu\text{l/well}$ blocking buffer without BSA) repeated three times after each step. The coating solution was discarded, and blocking buffer (TBS, 10 mM CaCl₂, 0.05% Tween 20, 2% BSA) was added. The blocking buffer was discarded, and the desired representative Xyn-Doc(s) (Xyn-DocA, Xyn-DocB, Xyn-DocC, Xyn-DocD, Xyn-DocF, Xyn-DocG, Xyn-DocH, Xyn-DocJ, Xyn-Doc3939, Xyn-Doc48, Xyn-Doc9b, Xyn-Doc3550), diluted to concentrations of 0.1, 1, 10 and 100 ng ml^{-1} in blocking buffer, were added. Rabbit anti-xylanase antibody (diluted 1:10 000) was used as the primary antibody preparation, and secondary antibody preparation was HRP-labeled anti-rabbit antibody diluted

1:10 000 in blocking buffer. Substrate-Chromogen TMB (Dako, Agilent Technologies, USA) was added at 100 $\mu\text{l/well}$, and the reaction was carried out for 2 min before color formation was terminated upon addition of 1 M H₂SO₄ (50 $\mu\text{l/well}$), and the absorbance was measured at 450 nm using a tunable microplate reader.

Enzymatic activity assay

All assays were performed at least twice in triplicate. ScaK enzymatic activity was monitored using the turbidimetric method (Shugar, 1952; Diez-Martinez *et al.*, 2015) with modifications. The enzyme was applied at a concentration of 50 $\mu\text{g ml}^{-1}$, and *E. coli* BL21 at $\text{OD}_{600 \text{ nm}} = 1$ was used as a substrate for examining pH and temperature effects on the enzyme. After 30 min incubation time, decreases in optical densities were read at 600 nm after centrifuging 2 min at 1000 rpm ($60 \times g$) to clear cell debris. At this centrifugal speed, less than 10% of the bacterial cells precipitate, for short time periods (2 min) as can be observed in Supporting Information Fig. S5A and B. Without the centrifugation step, cell debris (that precipitate only gradually over extended time periods) can float and OD measurements can be less accurate. Controls without enzyme were also submitted to the centrifugal step so that the 10% precipitation of the bacterial cells does not affect the results. Supporting Information Fig. S5C presents the effect of temperature on lytic activity after 1 h incubation without the centrifugation step, and the results tend to be similar to those presented in Fig. 4C with the centrifugation step.

The pH optimum was determined by using acetate buffer ranging from pH 3.5 to 6.5 and MOPS (3-(*N*-morpholino)propanesulfonic acid) from pH 6.5 to 9. The enzyme was incubated at 37°C for 15 min. Temperature optima were tested at pH 5 (50 mM acetate buffer) between temperatures ranging from 2 to 45°C, and reactions were terminated after 30 min incubation. We used a viability assay (Diez-Martinez *et al.*, 2015) to follow the temperature effect on ScaK lytic action. Measurement of viable *E. coli* cells was carried out in LB agar plates after 30 min incubation at each temperature. For each sample, a 10-fold dilution series was prepared in LB, and 100 μl of each dilution was plated. Colonies were counted after overnight incubation at 37°C.

The susceptibility of the peptidoglycans from *B. subtilis* and *S. aureus* (Sigma-Aldrich, Israel) to lysozyme was analysed with a turbidometric assay (Bera *et al.*, 2005; Wang *et al.*, 2009). The peptidoglycans were diluted to 0.5 mg ml^{-1} in double distilled water, and the lysozyme was added to a concentration of 50 $\mu\text{g ml}^{-1}$ at pH 5 (50 mM acetate buffer). The absorbance at 450 nm was monitored after 30 min of incubation at temperatures ranging from 2 to 85°C.

Stability of the protein was tested for 30 min at 37°C after incubation for 1, 3, 24 and 48 h at 30, 40, 50, 60, 70 and 80°C.

Bacterial strains and growth medium

The bacterial species tested in this study are listed in Table 2. All the strains were cultured at 37°C (anaerobic strains were grown under anaerobic conditions) and tested for their sensitivity to ScaK *in vitro* as described above (at 1 h incubation

time). Certain strains listed in Table 2 were held in stock by the authors S.H. Duncan and H.J. Flint (U. of Aberdeen). The 18 cultures provided for testing for ScaK lytic activity were *Bifidobacterium adolescentis* L2-32, *Collinsella aerofaciens* DSM 3979, *Enterococcus faecalis* JH2-2, *Eubacterium hallii* L2-7 (DSM 17630), *Anaerostipes hadrus* Ss2/1, *Blautia obeum* A2-162, *Eubacterium rectale* A1-86 (DSM 17629), *Roseburia faecis* M72/1 (DSM 16840), *Lachnospiraceae* sp. nov. M62/1, *Lactobacillus reuteri* Ca6, *R. champanellensis* 18P13 (DSM 18848), *Ruminococcus bicirculans* 80/3, *Ruminococcus bromii* L2-63, *Eubacterium siraeum* 70/3, *Faecalibacterium prausnitzii* A2-165 (DSM 17677), *Ruminococcus flavefaciens* 17, *Ruminococcus albus* SY3 and *Streptococcus gordonii* DL-1. Those available from Deutsche Sammlung von Mikroorganismen und Zellkulturen (DSMZ), Germany) are indicated by DSM numbers in brackets. Cultures were prepared by growing on M2GSC medium (Miyazaki *et al.*, 1997) for approximately 24 h under CO₂ prior to testing for ScaK lysis. *E. coli* BL21 and Nissle, *Bacillus subtilis* NCIB 3610 and PY79 strains, *Acinetobacter baumannii* ATCC 17978, *Listeria monocytogenes* 10403S and *Vibrio cholerae* TRH7000 were cultured in LB. *Bacteroides thetaiotaomicron* ATCC 29148 and ΔCPS (Cameron *et al.*, 2014) were cultured in TYG (Holdeman *et al.*, 1977). *L. plantarum* WCFS1, *Lactobacillus pentosus* DSM 20314, *Lactobacillus reuteri* Ca6 and *Lactobacillus rhamnosus* GG were cultured in MRS. *Lactococcus lactis* MG5267 and Z3000 strains were cultured in M17. *Clostridium difficile* VPI 10463 was cultured in BHIS (Sorg and Dineen, 2009). *Clostridium thermocellum* DSM 1313 and *Clostridium clariflavum* DSM 19732 were cultured in GS-2 medium. *R. champanellensis* 18P13 was cultured in M2 medium. *S. pneumoniae* R6 was grown in Todd Hewitt broth supplemented with 5% yeast extract (Updyke and Nickle, 1954).

E. coli strain 5911 was grown for 3 h in LB, and capsule expression was induced by addition of 15% sucrose (Sledjeski and Gottesman, 1996). The cells were then grown overnight at 37°C and incubated at room temperature for a week.

Bacterial cells were tested by the turbidimetric assay as described above.

Inhibition test

E. coli BL21 and *L. plantarum* WCFS1 cells were spread on the entire surface of LB or MRS plates prepared at pH 5. Sterile filter paper disks (catalogue number 74146, Sigma Aldrich, Israel) containing either sterile water or 2 g l⁻¹ ScaK were placed in the middle of the plates, and the plates were incubated overnight at 37°C.

Label-free LC MS-MS analysis

Proteolysis and mass spectrophotometry analysis of *R. champanellensis* culture supernatants, grown on cellobiose as a carbon source (Lopez-Siles *et al.*, 2012; Morais *et al.*, 2016), were performed as described by Artzi *et al.* (2015).

Acknowledgements

This research was supported by a grant to EAB, NMK and ECM from the United States-Israel Binational Science

Foundation (BSF), Jerusalem, Israel and by a grant (No. 1349) to EAB from the Israel Science Foundation (ISF). Additional support was obtained from the establishment of an Israeli Center of Research Excellence (I-CORE Center No. 152/11) managed by the Israel Science Foundation. The authors also appreciate the support of the European Union, Area NMP2013.1.1-2: Self-assembly of naturally occurring nanosystems: CellulosomePlus Project number: 604530. HJF and SHD acknowledge financial support from the Scottish Government Rural and Environmental Sciences and Analytical Services (SG-RESAS) and from BBSRC grant no BB/L009951/1. EAB is the incumbent of The Maynard I. and Elaine Wishner Chair of Bio-organic Chemistry.

Conflict of interest

The authors declare no conflict of interest

References

- Alahuhta, M., Xu, Q., Brunecky, R., Adney, W.S., Ding, S.Y., Himmel, M.E., and Lunin, V.V. (2010) Structure of a fibronectin type III-like module from *Clostridium thermocellum*. *Acta Crystallogr Sect F Struct Biol Cryst Commun* **66**: 878–880.
- Aminlari, L., Hashemi, M.M., and Aminlari, M. (2014) Modified lysozymes as novel broad spectrum natural antimicrobial agents in foods. *J Food Sci* **79**: R1077–R1090.
- Artzi, L., Morag, E., Barak, Y., Lamed, R., and Bayer, E.A. (2015) *Clostridium clariflavum*: key cellulosome players are revealed by proteomic analysis. *mBio* **6**: e00411–e00415.
- Barak, Y., Handelsman, T., Nakar, D., Mechaly, A., Lamed, R., Shoham, Y., and Bayer, E.A. (2005) Matching fusion protein systems for affinity analysis of two interacting families of proteins: the cohesin–dockerin interaction. *J Mol Recognit* **18**: 491–501.
- Bayer, E.A., Belaich, J.P., Shoham, Y., and Lamed, R. (2004) The cellulosomes: multi-enzyme machines for degradation of plant cell wall polysaccharides. *Annu Rev Microbiol* **58**: 521–554.
- Bayer, E.A., Lamed, R., and Himmel, M.E. (2007) The potential of cellulases and cellulosomes for cellulosic waste management. *Curr Opin Biotechnol* **18**: 237–245.
- Ben David, Y., Dassa, B., Borovok, I., Lamed, R., Koropatkin, N.M., Martens, E.C., *et al.* (2015) Ruminococcal cellulosome systems from rumen to human. *Environ Microbiol* **17**: 3407–3426.
- Bera, A., Herbert, S., Jakob, A., Vollmer, W., and Gotz, F. (2005) Why are pathogenic staphylococci so lysozyme resistant? The peptidoglycan O-acetyltransferase OatA is the major determinant for lysozyme resistance of *Staphylococcus aureus*. *Mol Microbiol* **55**: 778–787.
- Berg Miller, M.E., Yeoman, C.J., Chia, N., Tringe, S.G., Angly, F.E., Edwards, R.A., *et al.* (2012) Phage-bacteria relationships and CRISPR elements revealed by a metagenomic survey of the rumen microbiome. *Environ Microbiol* **14**: 207–227.
- Berlemont, R., and Martiny, A.C. (2013) Phylogenetic distribution of potential cellulases in bacteria. *Appl Environ Microbiol* **79**: 1545–1554.

- Briers, Y., and Lavigne, R. (2015) Breaking barriers: expansion of the use of endolysins as novel antibacterials against Gram-negative bacteria. *Future Microbiol* **10**: 377–390.
- Callewaert, L., Walmagh, M., Michiels, C.W., and Lavigne, R. (2011) Food applications of bacterial cell wall hydrolases. *Curr Opin Biotechnol* **22**: 164–171.
- Callewaert, L., Van Herreweghe, J.M., Vanderkelen, L., Leysen, S., Voet, A., and Michiels, C.W. (2012) Guards of the great wall: bacterial lysozyme inhibitors. *Trends Microbiol* **20**: 501–510.
- Cameron, E.A., Kwiatkowski, K.J., Lee, B.H., Hamaker, B.R., Koropatkin, N.M., and Martens, E.C. (2014) Multifunctional nutrient-binding proteins adapt human symbiotic bacteria for glycan competition in the gut by separately promoting enhanced sensing and catalysis. *mBio* **5**: 01414–e01441.
- Caspi, J., Irwin, D., Lamed, R., Shoham, Y., Fierobe, H.P., Wilson, D.B., and Bayer, E.A. (2006) *Thermobifida fusca* family-6 cellulases as potential designer cellulosome components. *Biocatal Biotransform* **24**: 3–12.
- Chassard, C., Delmas, E., Robert, C., Lawson, P.A., and Bernalier-Donadille, A. (2012) *Ruminococcus champanellensis* sp. nov., a cellulose-degrading bacterium from human gut microbiota. *Int J Syst Evol Microbiol* **62**: 138–143.
- Claverys, J.P., Martin, B., and Havarstein, L.S. (2007) Competence-induced fratricide in streptococci. *Mol Microbiol* **64**: 1423–1433.
- Datta, S., Janes, M.E., Xue, Q.G., Losso, J., and La Peyre, J.F. (2008) Control of *Listeria monocytogenes* and *Salmonella anatum* on the surface of smoked salmon coated with calcium alginate coating containing oyster lysozyme and nisin. *J Food Sci* **73**: M67–M71.
- Devillard, E., Goodheart, D.B., Karnati, S.K., Bayer, E.A., Lamed, R., Miron, J., et al. (2004) *Ruminococcus albus* 8 mutants defective in cellulose degradation are deficient in two processive endocellulases, Cel48A and Cel9B, both of which possess a novel modular architecture. *J Bacteriol* **186**: 136–145.
- Diez-Martinez, R., De Paz, H.D., Garcia-Fernandez, E., Bustamante, N., Euler, C.W., Fischetti, V.A., et al. (2015) A novel chimeric phage lysin with high in vitro and in vivo bactericidal activity against *Streptococcus pneumoniae*. *J Antimicrob Chemother* **70**: 1763–1773.
- Eldholm, V., Johnsborg, O., Haugen, K., Ohnstad, H.S., and Havarstein, L.S. (2009) Fratricide in *Streptococcus pneumoniae*: contributions and role of the cell wall hydrolases CbpD, LytA and LytC. *Microbiology* **155**: 2223–2234.
- Fouet, A., and Mesnage, S. (2002) *Bacillus anthracis* cell envelope components. In *Anthrax*. Koehler, T.M. (ed). Berlin, Heidelberg: Springer Berlin Heidelberg, pp. 87–113.
- Gasteiger, E., Hoogland, C., Gattiker, A., Duvaud, S., Wilkins, M.R., Appel, R.D., and Bairoch, A. (2005) Protein identification and analysis tools on the ExPASy server. In *The Proteomics Protocols Handbook*. Walker, J.M. (ed). Totowa, NJ: Humana Press, pp. 571–607.
- Gottesman, S., and Stout, V. (1991) Regulation of capsular polysaccharide synthesis in *Escherichia coli* K12. *Mol Microbiol* **5**: 1599–1606.
- Himmel, M., Xu, Q., Luo, Y., Ding, S., Lamed, R., and Bayer, E. (2010) Microbial enzyme systems for biomass conversion: Emerging paradigms. *Biofuels* **1**: 323–341.
- Holdeman, L.V., Cato, E.D., and Moore, W.E.C. (1977) *Anaerobe Laboratory Manual*. Blacksburg, VA: Virginia Polytechnic Institute and State University Anaerobe Laboratory.
- Humann, J., and Lenz, L.L. (2009) Bacterial peptidoglycan degrading enzymes and their impact on host muropeptide detection. *J Innate Immunity* **1**: 88–97.
- Lai, M.J., Lin, N.T., Hu, A., Soo, P.C., Chen, L.K., Chen, L.H., and Chang, K.C. (2011) Antibacterial activity of *Acinetobacter baumannii* phage varphiAB2 endolysin (LysAB2) against both gram-positive and gram-negative bacteria. *Appl Microbiol Biotechnol* **90**: 529–539.
- Lohning, C., Pohlschmidt, M., Glucksmann-Kuis, M.A., Duyk, G., Bork, P., Schneider, M.O., et al. (1996) Structural motifs of the PKD1 protein. *Nephrol Dial Transplant* **11**(Suppl 6): 2–4.
- Lood, R., Winer, B.Y., Pelzek, A.J., Diez-Martinez, R., Thandar, M., Euler, C.W., et al. (2015) Novel phage lysin capable of killing the multidrug-resistant gram-negative bacterium *Acinetobacter baumannii* in a mouse bacteremia model. *Antimicrob Agents Chemother* **59**: 1983–1991.
- Lopez-Siles, M., Khan, T.M., Duncan, S.H., Harmsen, H.J., Garcia-Gil, L.J., and Flint, H.J. (2012) Cultured representatives of two major phylogroups of human colonic *Faecalibacterium prausnitzii* can utilize pectin, uronic acids, and host-derived substrates for growth. *Appl Environ Microbiol* **78**: 420–428.
- Ludwig, W., Schleifer, K.H., Whitman, W.B. (2009) Revised road map to the phylum Firmicutes. In *Bergey's Manual® of Systematic Bacteriology: Volume Three the Firmicutes*. Vos, P., Garrity, G.M., Jones, D., Krieg, N.R., Ludwig, W., Rainey, F.A., et al. (eds). New York, NY: Springer New York, pp. 1–13.
- Lytle, B., Myers, C., Kruus, K., and Wu, J.H.D. (1996) Interactions of the CelS binding ligand with various receptor domains of the *Clostridium thermocellum* cellulosomal scaffolding protein, CipA. *J Bacteriol* **178**: 1200–1203.
- Miyazaki, K., Martin, J.C., Marinsek-Logar, R., and Flint, H.J. (1997) Degradation and utilization of xylans by the rumen anaerobe *Prevotella bryantii* (formerly *P. ruminicola* subsp. *brevis*) B(1)4. *Anaerobe* **3**: 373–381.
- Monterroso, B., Saiz, J.L., Garcia, P., Garcia, J.L., and Menendez, M. (2008) Insights into the structure-function relationships of pneumococcal cell wall lysozymes, LytC and Cpl-1. *J Biol Chem* **283**: 28618–28628.
- Morais, S., David, Y.B., Bensoussan, L., Duncan, S.H., Koropatkin, N.M., Martens, E.C., et al. (2016) Enzymatic profiling of cellulosomal enzymes from the human gut bacterium, *Ruminococcus champanellensis*, reveals a fine-tuned system for cohesin-dockerin recognition. *Environ Microbiol* **18**: 542–556.
- Pastagia, M., Schuch, R., Fischetti, V.A., and Huang, D.B. (2013) Lysins: the arrival of pathogen-directed anti-infectives. *J Med Microbiol* **62**: 1506–1516.
- Rincon, M.T., Martin, J.C., Aurilia, V., McCrae, S.I., Rucklidge, G.J., Reid, M.D., et al. (2004) ScaC, an adaptor protein carrying a novel cohesin that expands the dockerin-binding repertoire of the *Ruminococcus flavefaciens* 17 cellulosome. *J Bacteriol* **186**: 2576–2585.
- Rogers, T.E., Pudlo, N.A., Koropatkin, N.M., Bell, J.S., Moya Balasch, M., Jasker, K., and Martens, E.C. (2013) Dynamic responses of *Bacteroides thetaiotaomicron* during growth on glycan mixtures. *Mol Microbiol* **88**: 876–890.

- Schleifer, K.H., and Kandler, O. (1972) Peptidoglycan types of bacterial cell walls and their taxonomic implications. *Bacteriol Rev* **36**: 407–477.
- Shah, H.N. (1992) The Genus *Bacteroides* and Related Taxa. In *The Prokaryotes: A Handbook on the Biology of Bacteria: Ecophysiology, Isolation, Identification, Applications*. Balows A., Trüper H.G., Dworkin M., Harder W., Schleifer K.-H. (eds), New York, NY: Springer New York, pp. 3593–3607.
- Shugar, D. (1952) The measurement of lysozyme activity and the ultra-violet inactivation of lysozyme. *Biochim Biophys Acta* **8**: 302–309.
- Sledjeski, D.D., and Gottesman, S. (1996) Osmotic shock induction of capsule synthesis in *Escherichia coli* K-12. *J Bacteriol* **178**: 1204–1206.
- Sorg, J.A., and Dineen, S.S. (2009) Laboratory maintenance of *Clostridium difficile*. *Curr Protoc Microbiol* Chapter 9:Unit9A 1.
- Torres-Cabassa, A.S., and Gottesman, S. (1987) Capsule synthesis in *Escherichia coli* K-12 is regulated by proteolysis. *J Bacteriol* **169**: 981–989.
- Updyke, E.L., and Nickle, M.I. (1954) A dehydrated medium for the preparation of type specific extracts of group A *Streptococci*. *Appl Microbiol* **2**: 117–118.
- Vollmer, W., Joris, B., Charlier, P., and Foster, S. (2008) Bacterial peptidoglycan (murein) hydrolases. *FEMS Microbiol Rev* **32**: 259–286.
- Wang, G., Olczak, A., Forsberg, L.S., and Maier, R.J. (2009) Oxidative stress-induced peptidoglycan deacetylase in *Helicobacter pylori*. *J Biol Chem* **284**: 6790–6800.
- Wegmann, U., Louis, P., Goesmann, A., Henrissat, B., Duncan, S.H., and Flint, H.J. (2014) Complete genome of a new Firmicutes species belonging to the dominant human colonic microbiota (*Ruminococcus bicirculans*) reveals two chromosomes and a selective capacity to utilize plant glucans. *Environ Microbiol* **16**: 2879–2890.
- Whitfield, C., and Roberts, I.S. (1999) Structure, assembly and regulation of expression of capsules in *Escherichia coli*. *Mol Microbiol* **31**: 1307–1319.
- Yaron, S., Morag, E., Bayer, E.A., Lamed, R., and Shoham, Y. (1995) Expression, purification and subunit-binding properties of cohesins 2 and 3 of the *Clostridium thermocellum* cellulosome. *FEBS Lett* **360**: 121–124.

Supporting information

Additional supporting information may be found in the online version of this article at the publisher's web-site:

Table S1. Analysis of the flanking regions of the ScaK gene.

Fig. S1. Purity of the recombinant scaffoldins after Ni-NTA purification as assessed by SDS-PAGE gels. A. ScaK, molecular weight 52326 Da (10% acrylamide), B. ScaG, molecular weight 29510 Da (12% acrylamide) and C. ScaF, molecular weight 25633 Da (12% acrylamide).

Fig. S2. *R. champanellensis* ScaF interactions measured by ELISA with selected dockerins. The cohesin only binds to the dockerin of ScaG. Error bars indicate the standard deviation from the mean of triplicate (ELISA) from three experiments.

Fig. S3. Proposed cell-bound cellulosome complexes involving ScaG and ScaF in *R. champanellensis*. Different types of cohesin-dockerin interactions are color-coded. The ScaG and ScaF dockerins are selective for the cohesins of ScaE and ScaJ1. The ScaG dockerin also binds selectively to the ScaF cohesin, but the ScaF cohesin fails to bind to its own dockerin. The ScaG cohesin appears to be very selective in its binding to the ScaA dockerin and fails to bind other Group 2 dockerins.

Fig. S4. Inhibition of *L. plantarum* (A) and *E. coli* (B) growth by the presence of the GH25 lysozyme on disks (2 g l^{-1}). The cells were grown on MRS and LB plates prepared at pH 5. Inhibition zones are marked with red arrows.

Fig. S5. Effect of time on the precipitation of bacterial cells *L. plantarum* (A) and *E. coli* (B) at a centrifugal speed of 1000 rpm ($60 \times g$). (C) Effect of temperature on lysozyme activity on *E. coli* cells after 1 h incubation at pH 5 without the centrifugation step.